

Homogenic Evaluation for Spatial Distribution in Osteoclast Differentiation

Hyun-Sook Lim¹, Hong-In Shin^{2,*}, Daewon Jeong^{3,4,*}

¹Department of Public Health Administration, Hanyang Women's University, Seoul;

²Department of Oral Pathology, School of Dentistry, Institute for Hard Tissue and Bio-Tooth Regeneration, Kyungpook National University, Daegu;

³Laboratory of Bone Metabolism and Control, Department of Microbiology, Yeungnam University College of Medicine, Daegu;

⁴Company of the Bone Science, Yeungnam University College of Medicine, Daegu, Korea

Corresponding authors

Hong-In Shin

Department of Oral Pathology, School of Dentistry, Institute for Hard Tissue and Bio-Tooth Regeneration, Kyungpook National University, 2177 Dalgubeol-daero, Jung-gu, Daegu 41940, Korea
Tel: +82-53-420-4994
Fax: +82-53-428-4995
E-mail: hishin@knu.ac.kr

Daewon Jeong

Laboratory of Bone Metabolism and Control, Department of Microbiology, Yeungnam University College of Medicine, 170 Hyeonchung-ro, Nam-gu, Daegu 42415, Korea
Tel: +82-53-640-6944
Fax: +82-53-657-6869
E-mail: dwjeong@ynu.ac.kr

Received: November 8, 2022

Revised: November 17, 2022

Accepted: November 21, 2022

*Hong-In Shin and Daewon Jeong contributed equally to this work and should be considered co-corresponding authors.

Copyright © 2022 The Korean Society for Bone and Mineral Research

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.




Background: Cells have heterogeneous cellular diversity in size, morphology, cell cycle, metabolism, differentiation degree, and spatial distribution. The shift of specific cells towards the desired cells is crucial for maintaining uniform cellular function and can be represented by homogeneity and heterogeneity. Here, we developed a simple and direct method for evaluating the homogeneous distribution of desired cells in a constant region. **Methods:** We differentiated osteoclast progenitors into bone-resorbing multinucleated giant osteoclasts in a 2-dimensional culture plate under 2 conditions. Cells were stained with tartrate-resistant acid phosphatase to assess osteoclast differentiation, images were taken using a microscope and divided into sectors, and the number of osteoclasts (≥ 3 nuclei) in each sector was counted. To assess the homogeneity of the spatial distribution of osteoclasts, the standard deviation (SD) was calculated from the mean number of osteoclasts within each sector. **Results:** From the 2 groups, a value with a SD close to 0 indicates high spatial homogeneity while a relatively high SD represents low spatial homogeneity. **Conclusions:** Our findings suggest that spatial homogeneity can be represented as SD.

Key Words: Cell differentiation · Homogeneity · Heterogeneity · Osteoclast · Spatial distribution

INTRODUCTION

Cells are necessary for switching between homogeneous and heterogeneous states according to circumstances. Unidirectional differentiation and proliferation of stem cells into a homogeneous cell population are effective in maintaining uniform cellular function.[1] A high yield of cells of interest by reducing heterogeneous cell populations is also important for improving cell therapy to treat diseases.[1,2] Conversely, the drift of homogeneous cells to heterogeneous cells confers specialized functions in physiological and pathological conditions. Heterogeneous cells derived from homogeneous cells are favorable for coping with changing environments [3]; on the contrary, cancer cell heterogeneity caused by genetic mis-regulations (e.g., genomic instability, epigenetic alteration, and plastic gene expression), signal transduction, and microenvironmental differences drives

Graphical Abstract



phenotypic diversity (heterogeneity), resulting in resistance to anticancer treatment).[4] Hereafter, the biotechnology-based development of heterogeneous cancer cells with resistance to anticancer treatment that can induce homogeneous cancer cells with sensitivity to anticancer drugs becomes an important issue.[5,6] Thus, a proper balance between heterogeneous and homogeneous cells is required to regulate cell capacity in various environments under time and space of physiological conditions.

In *in vitro* cell culture, cells show differences between neighboring cells in the same area and space. Their cells display differences in external cellular phenotypes (e.g., cell size and morphology) and internal cellular events that happen inside the cell (e.g., cell cycle, metabolic and cellular signaling pathways, and transcriptional and translational activities).[7] In addition, the final cells formed under specific conditions differ in their degree of differentiation and spatial distribution.[8] This difference could result from various factors, such as medium composition, cell dispersion and confluence, and heterogeneous cell populations with genetic background variation. Reprogramming heterogeneous cells to homogeneous cells will act favorably in inducing cells with a uniform function. For instance, mes-

enchymal stem cells with ectopic expression of specific transcription factors promote trans-differentiation into osteogenic, chondrogenic, adipogenic and myogenic cell lineage [9] or synchronized osteoclast progenitors with cell cycle arrest by macrophage colony-stimulating factor (M-CSF) depletion effectively undergo uniform osteoclast differentiation.[10] Among the cellular heterogenic diversities in microenvironments that vary in time and space, the spatial distribution of cells serves as a cell fate determinant of migration, differentiation, proliferation, and programmed cell death.[11-14] This is also a key player in tissue development.[15]

Parallel to the technical exploration of switching between heterogeneous and homogeneous cells, a method for evaluating homogeneity is required to accurately define the degree of homogeneity. In this study, we developed a simple method for assessing the extent of the spatial distribution of a certain cell in a constant area. The mean value of the desired cell number in a region was obtained from all cell numbers within the total area, and the standard deviation (SD) was then calculated from the deviated value of the mean. Small and large SDs represent high and low homogeneities, respectively. This method is

useful for assessing the spatial distribution of the targeted cells in 2-dimensional (D) area and 3D space.

METHODS

To generate osteoclast progenitors, bone marrow cells were flushed out from the femur and tibia of 6-week-old male C5BL/6J mice (Central Lab Animal, Seoul, Korea) using α -minimal essential medium (α -MEM; HyClone, Logan, UT, USA). Cells were plated on cell culture plates in α -MEM with 10% fetal bovine serum (FBS) and 5 ng/mL M-CSF, and incubated in a humidified incubator with 5% CO₂ at 37°C overnight. Non-adherent cells were harvested and cultured in α -MEM containing 10% FBS and 30 ng/mL M-CSF for 2 days to produce osteoclast progenitors. For osteoclast differentiation, osteoclast progenitors (1×10^4 cells/well or 2×10^4 cells/well in a 24-well culture plate) were cultured in α -MEM with M-CSF (30 ng/mL) and receptor activator of nuclear factor- κ B ligand (RANKL; 100 ng/mL) for 4 days, with fresh medium replaced after 2 days. To assess osteoclast formation, cells were washed with phosphate-buffered saline, fixed with 3.7% (v/v) formaldehyde, and stained for tartrate-resistant acid phosphatase (TRAP) using a commercial leukocyte acid phosphatase staining kit (Sigma-Aldrich, St. Louis, MO, USA). After cell images were taken under a light microscope, TRAP-positive multinucleated cells (TRAP+MNCs) with more than 3 nuclei were counted to assess the extent of osteoclast differentiation in a constant area.

RESULTS AND DISCUSSION

1. Three formulas for assessing cellular homogeneity in spatial distribution

To assess the spatial distribution, the images taken under different cell conditions were divided into sectors with a regular quadrangle. After counting specific cells within each sector and adding all cell numbers, the mean, a statistical average, was calculated by dividing the sum of all the numbers within the sector by the number of sectors, as described in equation (1).

$$m = \frac{\sum_{i=1}^n S_i}{n} \quad (1)$$

Where m is the average value (mean) for all cell numbers

of each sector, n is the sector number, and S_i is the cell number in each sector.

Next, to characterize the scatter around the mean value, the variance (V) of a sector n , defined in equation (2), was calculated.

$$V = \frac{\sum_{i=1}^n (S_i - m)^2}{n} \quad (2)$$

Finally, the SD (σ), which is a measure of the amount of variation or dispersion of a set of values, was calculated from the square root of the variance [as described in equation (3)]. [16]

$$\sigma = \sqrt{V} \quad (3)$$

A low SD indicates that the values are clustered closely around the mean, whereas a high SD indicates that they are scattered far from the mean. Thus, the SD deduced from the resulting mean can be applied to evaluate the spatial distribution of the targeted cells in a constant region.

2. Application of SD to homogenic assessment for the spatial distribution of osteoclast differentiation

To demonstrate the possibility that SD can be applied to assess the spatial homogeneity of a certain cell, we attempted to induce the differentiation of bone marrow-derived osteoclast progenitors into bone-resorbing multinucleated osteoclasts. As shown in Figure 1, osteoclast progenitors with cell numbers of 1×10^4 cells/well (group A) and 2×10^4 cells/well (group B) in 24-well plates were cultured in a medium containing M-CSF. When the cells reached approximately 40% (group A) and 60% confluence (group B), they were differentiated into osteoclasts in a conditioned medium containing M-CSF and RANKL for 4 days. To assess the homogeneous distribution of differentiated osteoclasts, cell images were taken from the 2 groups (A and B) under a microscope. After the images were divided into regular quadrangle sections, osteoclasts with more than 3 nuclei within each sector were counted. The mean was obtained by dividing the sum of the cell numbers for all sectors by the number of sectors, and the SD was estimated from the resulting mean. SD provides the extent of homogeneous distribution between groups, whether the values converge to and diverge from the mean. The SDs for groups A and B were 1.946 and 1.138, respectively, indicating that the osteoclasts in group B, which had a lower SD, were more evenly distributed in the 2D culture condition than those in




Fig. 1. Evaluation for homogenic distribution of differentiated osteoclasts in 2-dimensional culture condition. Osteoclast progenitors of 1×10^4 cells/well (group A) and 2×10^4 cells/well (group B) were plated on a 24-well culture plate and cultured in a medium with macrophage colony-stimulating factor (M-CSF) until cell confluence reached approximately 40% (group A) and 60% (group B). Cells were further differentiated into osteoclasts in a medium containing M-CSF and receptor activator of nuclear factor- κ B ligand for 4 days with the medium changed after 2 days. Cells formed under different conditions were stained for tartrate-resistant acid phosphatase (TRAP) to detect osteoclasts and were imaged under a light microscope at the same magnification (step 1). The images were divided into sectors and TRAP-positive osteoclasts with more than 3 nuclei within each sector in group A and group B were counted (step 2). The extent of homogenic distribution of osteoclasts in a constant field was assessed through serial calculation of mean (m), variance (V), and standard deviation (σ) (scale bar, 100 μ m).




Fig. 2. Graphic illustration for the spatial distribution of differentiated osteoclasts. The values of mean and standard deviation (SD) obtained from Figure 1 were illustrated as a graph to visualize the data.

group A, which had a higher SD. Moreover, the graphs of the mean and SD can be used to visualize the extent of homogeneous distribution (Fig. 2). The higher homogeneity in the distribution of osteoclasts in group B compared to those in group A could be attributed to the fine-tuned communication between adjacent mononuclear osteoclast progenitors and the effective sequential processes via increased mononuclear cell migration and fusion.

In addition to technical advances that can control the switching between homogeneous and heterogeneous cells, a method capable of determining homogeneity is emerging for achieving a homogeneous assessment. Here,

we suggest a useful method for evaluating the homogeneous distribution of specific cells in a 2D area and 3D geometry.

DECLARATIONS

Funding

This research was funded by the National Research Foundation of Korea, grant numbers NRF-2022R1A2C1002736 and NRF-2022R1A5A2018865.

Conflict of interest

No potential conflict of interest relevant to this article was reported.

ORCID

Hyun-Sook Lim <https://orcid.org/0000-0003-4321-6963>

Hong-In Shin <https://orcid.org/0000-0001-7758-5826>

Daewon Jeong <https://orcid.org/0000-0001-8158-4279>

REFERENCES

1. Muller-Sieburg CE, Sieburg HB, Bernitz JM, et al. Stem cell heterogeneity: implications for aging and regenerative medicine. *Blood* 2012;119:3900-7. <https://doi.org/10.1182/blood-2011-12-376749>.

2. Cho CH, Parashurama N, Park EY, et al. Homogeneous differentiation of hepatocyte-like cells from embryonic stem cells: applications for the treatment of liver failure. *FASEB J* 2008;22:898-909. <https://doi.org/10.1096/fj.06-7764com>.
3. Donati G, Watt FM. Stem cell heterogeneity and plasticity in epithelia. *Cell Stem Cell* 2015;16:465-76. <https://doi.org/10.1016/j.stem.2015.04.014>.
4. Sun XX, Yu Q. Intra-tumor heterogeneity of cancer cells and its implications for cancer treatment. *Acta Pharmacol Sin* 2015;36:1219-27. <https://doi.org/10.1038/aps.2015.92>.
5. Marusyk A, Janiszewska M, Polyak K. Intratumor heterogeneity: The rosetta stone of therapy resistance. *Cancer Cell* 2020;37:471-84. <https://doi.org/10.1016/j.ccell.2020.03.007>.
6. Mansoori B, Mohammadi A, Davudian S, et al. The different mechanisms of cancer drug resistance: A brief review. *Adv Pharm Bull* 2017;7:339-48. <https://doi.org/10.15171/apb.2017.041>.
7. Lloyd AC. The regulation of cell size. *Cell* 2013;154:1194-205. <https://doi.org/10.1016/j.cell.2013.08.053>.
8. Ruiz SA, Chen CS. Emergence of patterned stem cell differentiation within multicellular structures. *Stem Cells* 2008;26:2921-7. <https://doi.org/10.1634/stemcells.2008-0432>.
9. Almalki SG, Agrawal DK. Key transcription factors in the differentiation of mesenchymal stem cells. *Differentiation* 2016;92:41-51. <https://doi.org/10.1016/j.diff.2016.02.005>.
10. Kwon M, Kim JM, Lee K, et al. Synchronized cell cycle arrest promotes osteoclast differentiation. *Int J Mol Sci* 2016;17. <https://doi.org/10.3390/ijms17081292>.
11. Gartner ZJ, Bertozzi CR. Programmed assembly of 3-dimensional microtissues with defined cellular connectivity. *Proc Natl Acad Sci U S A* 2009;106:4606-10. <https://doi.org/10.1073/pnas.0900717106>.
12. Nelson CM, Jean RP, Tan JL, et al. Emergent patterns of growth controlled by multicellular form and mechanics. *Proc Natl Acad Sci U S A* 2005;102:11594-9. <https://doi.org/10.1073/pnas.0502575102>.
13. Tekin H, Sanchez JG, Landeros C, et al. Controlling spatial organization of multiple cell types in defined 3D geometries. *Adv Mater* 2012;24:5543-7, 5542. <https://doi.org/10.1002/adma.201201805>.
14. Hui EE, Bhatia SN. Micromechanical control of cell-cell interactions. *Proc Natl Acad Sci U S A* 2007;104:5722-6. <https://doi.org/10.1073/pnas.0608660104>.
15. Chen CS, Mrksich M, Huang S, et al. Geometric control of cell life and death. *Science* 1997;276:1425-8. <https://doi.org/10.1126/science.276.5317.1425>.
16. Bland JM, Altman DG. Measurement error. *BMJ* 1996;312:1654. <https://doi.org/10.1136/bmj.312.7047.1654>.

